

# Investigation of Fermi-LAT Unassociated Sources in Milagro Emission Regions

Raspberry SIMPSON Advisers: Ester ALIU and Prof. Reshmi MUKHERJEE


### Outline

- Project Summary and Goals
- Project Context Gamma Ray Experimentation
- Research Methods and Analysis
- Conclusions and Future Work








Investigation of Fermi-LAT
Unassociated Sources in
Milagro Emission Regions




# Project Context









# gamma-rays



Milagro Sky Map



IACT Sky Map



Fermi 2FGL Map

The gamma ray band is so broad, that many instruments are needed to assemble the "full picture"

Energy


Milagro VERITAS Fermi - LAT

#### Milagro 1 TeV - 100 TeV

Milagro was a ground based water Cherenkov detector.








#### VERITAS 50 GeV - 50 TeV

VERITAS is a ground based imaging atmospheric Cherenkov detector.








#### Fermi - LAT 20 MeV - 300 MeV

Fermi is an imaging telescope located on a spacecraft at low earth orbit.








### VERITAS 50 GeV - 50 TeV

RITAS is a ground based aging atmospheric erenkov detector.








### Fermi - LAT 20 MeV - 300 MeV

Fermi is an imaging telescope located on a spacecraft at low earth orbit.








#### Milagro 1 TeV - 100 TeV

Milagro was a ground based water Cherenkov detector.








#### VERITAS 50 GeV - 50 TeV

VERITAS is a ground based imaging atmospheric Cherenkov detector.








### Fermi - LAT 20 MeV - 300 MeV

Fermi is an imaging telescoplocated on a spacecraft at lovearth orbit.








### Milagro 1 TeV - 100 TeV

Milagro was a ground based water Cherenkov detector.








### VERITAS 50 GeV - 50 7

VERITAS is a ground imaging atmospher Cherenkov detector







## Fermi 2FGL Catalog

Over 30 % of Fermi Sources are unidentified because of

- Poor resolution of instruments that make source localization difficult
- · Low photon statistics

The rest are mostly pulsars (5%) and blazars (45 %)

#### Blazars

Class of AGN (active galactic nuclei)

- · Highly variable
- Bright radio, X-ray and gamma ray emission



Artist's representation of a blazar

#### **Pulsars**

A Pulsating Star

- Rotating neutron star, emits beam of continuous radiation.
- Gamma-ray telescopes usually detect young, nearby pulsars




Crab - pulsar wind nebula

#### The rest are mostly pulsars (5%) and blazars (45 %)

#### Blazars

Class of AGN (active galactic nuclei)

- Highly variable
- Bright radio, X-ray and gamma ray emission



Artist's representation of a blazar

#### Pulsars

A Pulsating Star

- Rotating neutron star, emits beam of continuous radiation.
- Gamma-ray telescopes usually detect young, nearby pulsars




Crab - pulsar wind nebula


# Characterization of Fermi Sources

The Fermi experiment has found that two parameters that can help distinguish between pulsars and blazars are:


- variability index "defined as the  $\chi 2$  of the deviations of eleven monthly (30-day) source flux measurements from the average source flux"
- curvature significance "measures the deviation of the spectrum from the simple power-law form for each source."




# um from the simple power-law form for eac.




# Selection of Unassociated Sources








After making our selection, we found a total of 46 unassociated sources in the Milagro emission region.





# Milagro Sky Map of Galactic Plane













After making our selection, we found a total of 46 unassociated sources in the Milagro emission region.



# Statistical Approach -Logistic Regression

Gamma ray astrophysicists often rely on statistical methods to establish the significance of a data sample.

Following in that tradition, we used the statistical method of Logistic Regression to classify our Fermi sources as pulsars or blazars.




#### Logistic Regression - An Introduction

Logistic Regression: A statistical classification method that is used to predict binary outcomes.


Example\*: Suppose we wanted to understand the features that influence whether a political candidate wins or loses. Some possible parameters to consider may be the amount of money spent on a campaign or the amount of time spent campaigning.

Given a training set, we can determine our parameters and then use the hypothesis — function above to predict our outcomes.

$$\int_{\Theta} (x) = \frac{1 + \delta_{-\Theta_1} x}{1}$$




Once the computer learns how to draw this decision boundary, we can apply the algorithm to predict the win/lose outcome for a different data set.




# LR Model Implementation

We utilized python to implement our logistic regression model without the use of library packages.

Our training set included all known blazars and pulsars in the 2FGL catalog.







- · Address under-fitting of data
- · Add new parameters to model
- · Improve optimization method

# III uie zful catalog.

#### Variability vs. Curvature



#### Variability vs. Curvature



- Address under-fitting of data
- Add new parameters to model
- Improve optimization method

# Multi-wavelength Analysis

Investigate possible associations by determining if counterparts existed in other energy bands, i.e. X-ray.




SWIFT experiment



Chandra




XMM Newton



Suzaku



**SWIFT Images** 



# Multi-wavelength Analysis

Investigate possible associations by determining if counterparts existed in other energy bands, i.e. X-ray.




**SWIFT** experiment



Chandra



XMM Newton



Suzaku



**SWIFT Images** 















Fermi Source Name	Chandra	Swift-XRT	XMM-Newton
2FGL J1857.2+0055c	Exposure: 10070 s Offset: 5.9968 arcmin	Exposure: 453.215 s Offset: 6.1534 arcmin	None
2FGL J1842.8-0359c	None	Exposure: 513.23 s Offset: 8.4557 arcmin	Exposure: 26953 s Offset: 1.3321 arcmin
2FGL J1844.3-0343c	None	Exposure: 67904.679 s Offset: 1.0426 - 9.8324 arcmin	None
2FGL J1843.7-0312c	None	Exposure: 506.5 s Offset: 7.7901 arcmin	Exposure: 31919 s Offset: 9.5120 arcmin
2FGL J1847.2-0236	None	Exposure: 897.599 s Offset: 4.8058 arcmin	None
2FGL J1848.2-0139c	None	Exposure:493.93100 s Offset:6.8104 arcmin	Exposure: 65149 s Offset: 3.8670 arcmin
2FGL J1849.9-0125c	None	Exposure: 455.81900 s Offset: 4.3941 arcmin	None
2FGL J1857.2+0055c	None	Exposure: 453.21500 s Offset: 6.1534 arcmin	None
2FGL J1858.5+0129c	None	Exposure: 668.19800 s Offset: 5.9348 arcmin	None
2FGL J1857.6+0211	None	Exposure: 11127.638 s Offset: 1.5763 - 8.3669arcmin	None
2FGL J1859.3+0312c	None	Exposure: 2799.617 s Offset: 8.0175 - 9.4414 arcmin	None
2FGL J1857.8+0355c	None	Exposure: 493.027 s Offset: 5.3467 arcmin	None
2FGL J1901.1+0427	None	Exposure: 4715.848 s Offset: 0.362 - 9.4495 arcmin	None
2FGL J1906.5+0720	None	Exposure: 11264.895 s Offset: 1.0935 - 7.0505 arcmin	None
2FGL J1914.4+0951c	None	Exposure: 13069.055 s Offset: 0.8656 - 7.3889 arcmin	None
2FGL J1921.1+1436c	None	Exposure: 421.34600 s Offset: 6.8801 arcmin	None
2FGL J1914.0+1436	None	Exposure: 3756.999 s Offset: 1.4945 - 2.5999 arcmin	None
2FGL J1924.8+1724c	None	Exposure: 656.03900 s Offset: 3.6831 arcmin	None
2FGL J2028.3+3332	None	Exposure: 10427.484 s Offset: 1.3277 - 2.2050 arcmin	None
ABOT 100044 0000	**	Exposure: 4092.754 s	**


# SWIFT Images




















#### Conclusions

A combination quantitative study, the logistic regression model and qualitative study, the multi-wavelength comparisons, allow for promising investigation of Fermi's UFOs.

#### For the future ...

- Repeat LR model with Fermi 4th year catalog
- · Research other classification methods
- An expanded multi-wavelength study may help make improved associations.
- Complete XMM-Newton observation proposal (~ October)
- Make proposals for more observation of sources with VERITAS

promising investigation of Fermi's UFOs.

#### For the future ...

- Repeat LR model with Fermi 4th year catalog
- Research other classification methods
- An expanded multi-wavelength study may help make improved associations.
- Complete XMM-Newton observation proposal (~ October)
- Make proposals for more observation of sources with VERITAS

I would like to thank the Nevis REU program for giving the opportunity to work with my great advisers Ester Aliu, Reshmi Mukherjee and Gunes Senturk!

