

A brief, incomplete, and possibly incorrect introduction to statistics  
or  
A guide to physicists' jargon

The terms I hope to define


- Gaussian
- chi-squared
- chi-squared per degrees of freedom
- sigma (as in “we’re looking for a five-sigma effect”)
- systematic error

Professor Michael Shaevitz, Director of Nevis Labs, is our expert on statistics. I’m half-remembering what he taught me, and making up the rest.

# Gaussian

$$f(x) = e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

- $\mu$  is the mean of the distribution.
- $\sigma$  = the standard deviation; it's related to the “full width at half maximum” (FWHM) of the curve by  $\text{FWHM} = 2\sqrt{2\ln 2}\sigma \approx 2.35\sigma$ .
- $e$  = Euler's constant, a transcendental number that occurs often in calculations that relate to growth and increase. It's formally defined as  $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n$ .



## caveats

- If you want to work with the normal distribution as a “probability density function” then you’ll want to include a normalization so the integral  $\int_{-\infty}^{\infty} \mathcal{N}(x) dx = 1$


$$N(x; \mu, \sigma) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

However, if you work with this form you can’t fiddle with the amplitude of the distribution.

- There are other functional forms in physics than the Gaussian! However, I’m lazy, so that’s the only one I use in the tutorial.
- It makes some sense to stick with Gaussians, since the sum of many random processes (even non-Gaussian ones) tends towards a Gaussian.

# chi-squared

Measurements from my experiment



What's the probability that the underlying distribution of this histogram is a Gaussian?

## chi-square for a 1D histogram

$$\chi^2 = \sum_i \frac{(y_i - f(x_i; p_j))^2}{e_i^2}$$


where:

- $i$  means the  $i$ -th bin of the histogram (more generally, the  $i$ -th data point you've gathered).
- $y_i$  means the data (or value of) the  $i$ -th bin of the histogram.
- $e_i$  means the error in the  $i$ -th bin of the histogram (i.e., the size of the error bars).
- $f(x_i; p_j)$  means to compute the value of the function at  $x_i$  (the value on the  $x$ -axis of the center of bin  $i$ ) given some assumed values of the parameters  $p_0, p_1, p_2 \dots p_j$ .

The process of “fitting” means to test different values of the parameters until you find those that minimize the value of  $\chi^2$ .

Fit  $\equiv$  Test different values of  $A$ ,  $\mu$ , and  $\sigma$  until you find those that minimize the value of  $\chi^2$ .

Measurements from my experiment




The underlying program that does this is Minuit. It's been a standard program for finding function minima for decades.

# chi-squared per (number of) degrees of freedom

experiment

hist	
Entries	5000
Mean	5.015
Std Dev	1.932
$\chi^2 / \text{ndf}$	94.56 / 97
Constant	$99.53 \pm 1.80$
Mean	$5.025 \pm 0.030$
Sigma	$1.99 \pm 0.02$



## What value of chi-square do you expect from a fit?

$$\chi^2 = \sum_i \frac{(y_i - f(x_i; p_j))^2}{e_i^2}$$

- For each individual bin  $i$  the data forms a little gaussian distribution of its own with a mean of  $y_i$ .
- The  $e_i$  acts as a scale of the difference between  $y_i$  and the function  $f(x)$ . So if  $f(x)$  is a reasonable approximation to  $y_i$ ,  $(y_i - f(x))/e_i$  will be around  $\pm 1$ .
- You add up those “1”s for each of the bins, and you might anticipate that  $\chi^2$  will be roughly equal to  $i$ , the number of bins.

## But we have to adjust that...

There are three “free parameters” in the fit:  $A, \mu, \sigma$ .

They’re going to be varied to make the chi-squared smaller. The net effect is that total number of “degrees of freedom” is:

$$\begin{aligned} \text{DOF} &= \text{number of data points} \\ &\quad - \text{number of free parameters in the function} \end{aligned}$$

In that fit a couple of pages ago,  $\chi^2 / \text{ndf} = 94.56 / 97$ .

## What does this tell us?

Look it up on  
the web


**Upper-tail critical values of chi-square distribution with  $\nu$  degrees of freedom**

$\nu$	Probability less than the critical value				
	0.90	0.95	0.975	0.99	0.999
1	2.706	3.841	5.024	6.635	10.828
2	4.605	5.991	7.378	9.210	13.816
3	6.251	7.815	9.348	11.345	16.266
4	7.779	9.488	11.143	13.277	18.467
5	9.236	11.070	12.833	15.086	20.515
6	10.645	12.592	14.449	16.812	22.458

For the typical fits that we do in physics (lots of data points), it's sufficient that  $\chi^2 / \text{ndf}$  is roughly 1.

## Why might $\chi^2 / \text{ndf}$ be much greater than 1?


- There's something wrong in the routine that's calculating  $\chi^2$
- The model that's being assumed for the function does not have enough parameters.
- The error bars for your data are too small
- Function-minimization programs can get “stuck” in a local minimum that's not the actual true minimum



## Why might $\chi^2 / \text{ndf}$ be much less than 1?

- Again, something wrong in the  $\chi^2$  calculation.
- Too many free parameters in the function you're using to fit.
- The errors on your data are too large.
- Someone has gone wrong in your data-analysis process and you're “tuning” the data to the model you want to fit.


“We’re looking for a five sigma effect”



$3\sigma$  = “evidence”

$5\sigma$  = “discovery”


# This how I think of it



## Systematic error versus statistical error

Statistical error = random errors  
in the measurement.

Systematic error = a bias in the  
measurement, but you don't  
know how much that bias is.



## Example of an experiment's systematic errors

**ATLAS PUB Note**  
ATL-PHYS-PUB-2018-001  
31st January 2018

*Investigation of systematic  
uncertainties on the measurement  
of the top-quark mass using  
lepton transverse momenta*

Uncertainty	$\Delta m_{\text{top}}$ [GeV]
Statistics	0.94
Method calibration	0.40
Signal MC generator	0.62
Single-top Wt generator	0.28
Hadronisation and parton shower	0.55
ISR and FSR	1.39
Underlying Event	0.67
Colour Reconnection	0.23
Parton distribution function	0.42
Single-top contribution	0.10
Leptons	0.50
$E_{\text{T}}^{\text{miss}}$	0.12
$b$ -tagging	0.08
Jet energy scale	0.60
Jet energy resolution	0.32
Jet vertex fraction	0.05
Total	2.27

## Typical physicist lunchroom talk

- “What’s the chi-squared?” = What is the  $\chi^2$  per number of degrees of freedom from the fit to an assumed model (presumably a Gaussian)?
- “They’ve got a five-sigma effect!” = When you consider both the statistical and systematic errors, the measurement from the experiment refutes the null hypothesis at a large level of significance.
- “This sandwich tastes terrible!” = Let’s pick a different place to go to lunch next time.